

Contents

Part I

Preface	v
Sponsors and Supporters	vii
Committees	ix
Opening Address <i>Hiroshi Kamimura</i>	<i>xi</i>
Word of Welcome <i>Manuel Cardona</i>	<i>xiii</i>
Welcome Address <i>Hiroyuki Yoshikawa</i>	<i>xv</i>
Gracious Speech <i>Prince Takamado</i>	<i>xvii</i>
Telegram Message <i>Yoshiro Mori</i>	<i>xix</i>

1. Plenary Papers

Twenty years of quantum Hall effect <i>K. von Klitzing</i>	<i>1</i>
Optical manipulation of electron and nuclear spins in semiconductors <i>D.D. Awschalom</i>	<i>2</i>
Controlling the conductivity of wide-band-gap semiconductors <i>C.G. Van de Walle and J. Neugebauer</i>	<i>3</i>
High performance quantum cascade lasers for the mid-to far-infrared : from Band structure engineering to commercialization <i>F. Capasso</i>	<i>9</i>
Heterostructure lasers: development of new physics and new technology <i>Zh.I Alferov</i>	<i>14</i>
Quantum computation <i>Y. Yamamoto and F. Yamaguchi</i>	<i>20</i>
Hybrid structures of fullerenes and single-wall carbon nanotubes <i>S. Iijima</i>	<i>24</i>
The science, technology and impact of gallium nitride-based transistors <i>U.K. Mishra</i>	<i>25</i>
Correlation, decoherence, dephasing and relaxation in semiconductors <i>D.S. Chemla</i>	<i>26</i>
Experimental signatures of broken symmetries in the quantum Hall regime <i>J.P. Eisenstein</i>	<i>34</i>

2. Bulk and Dynamics

2.1 Electronic Structure

New effective-mass theory for degenerate bands in semiconductors <i>B.A. Foreman</i>	41
Antiferromagnetic coupling between the conduction electrons and the $4f$ electrons in $\text{Eu}@C_{60}$ <i>S. Suzuki, M. Kushida, S. Amamiya, S. Okada, and K. Nakao</i>	43
First-principles approach to the oxygen vacancies in SrTiO_3 <i>T. Shimizu and T. Kawakubo</i>	45
Hartree-Fock study on the phase diagrams of alkali-metal-doped C_{60} <i>J. Hirose, S. Suzuki, and K. Nakao</i>	47
Dynamical-mean-field study on the photoemission spectra of alkali-metal-doped C_{60} <i>T. Chida, S. Suzuki, and K. Nakao</i>	49
Spin effects in HgSe in megagauss fields <i>M. von Ortenberg, I. Stolpe, O. Portugall, N. Puhlmann, H.-U. Mueller, M. von Truchsess, C.R. Becker, A. Pfeuffer-Jeschke, and G. Landwehr</i>	51
Temperature dependence of effective band edge mass in $n\text{-InSb}$ by magnetophonon resonance up to $T = 320$ K and influence of nonparabolicity corrections <i>D. Schneider, S. Krull, C. Brink, G. Irmer, and P. Verma</i>	53
Self-energy correction to the mass of the wide gap semiconductors <i>M. Oshikiri and F. Aryasetiawan</i>	55
First-principles study of electronic structure of $\text{Si}_{1-x}\text{Ge}_x$ and $\text{Si}_{1-x-y}\text{Ge}_x\text{C}_y$ disordered alloys <i>M. Ohfuti, Y. Sugiyama, Y. Awano, and N. Yokoyama</i>	57
Electronic structures of the C_{82} and $\text{La}@C_{82}$ crystals by the relativistic LCAO method <i>S. Amamiya, S. Okada, S. Suzuki, and K. Nakao</i>	59
New determination of the camel's back in AlAs <i>L.E. Bremme, H. Im, H. Choi, P.C. Klipstein, R. Grey, and G. Hill</i>	61
Theory of the electronic structure of $\text{Ga}_{1-y}\text{In}_y\text{N}_x\text{As}_{1-x}$ <i>A. Lindsay and E.P. O'Reilly</i>	63
Influence of Ag on the energy spectrum of $(\text{Bi}_{1-x}\text{Sb}_x)_2\text{Te}_3$ <i>V.A. Kulbachinskii, A.Yu. Kaminsky, P. Lostak, and C. Drasar</i>	65

2.2 Phonons

Raman study of isotopically tailored CuBr <i>J. Serrano, F. Widulle, T. Ruf, C.T. Lin, and M. Cardona</i>	67
Isotope disorder effects in the Raman spectrum of Silicon <i>F. Widulle, T. Ruf, M. Konuma, I. Silier, W. Kriegseis, V.I. Ozhogin, and M. Cardona</i>	69
Observation of extraordinary phonon polaritons in ZnO <i>T. Azuhata, K. Torii, M. Ono, and T. Sota</i>	71
Disorder-activated resonant Raman scattering in GaNAs/GaAs structures <i>I.A. Buyanova, W.M. Chen, H.P. Xin, and C.W. Tu</i>	73
Collective dynamics in liquid Ge obtained by an inelastic x-ray scattering experiment <i>S. Hosokawa, Y. Kawakita, W.-C. Pilgrim, F. Hensel, H. Sinn, and M. Krisch</i>	75

2.3 Optical Properties and Excitons

Non-perturbative terahertz sideband generation from bulk GaAs <i>M.A. Zudov, J. Kono, A.P. Mitchell, A.H. Chin, and K. Johnsen</i>	77
--	----

Ab initio study of optical absorption spectra of semiconductors and conjugated polymers <i>M.L. Tiago, E.K. Chang, M. Rohlfing, and S.G. Louie</i>	79
Optical response of incoherent excitons in a dense electron-hole plasma <i>T. Schmielau, G. Manzke, D. Tamme, and K. Henneberger</i>	81
Effect of reabsorption on electroluminescence of excitonic origin in Cu ₂ O <i>Y. Nakamura, N. Naka, and N. Nagasawa</i>	83
Null Fresnel drag in a resonant moving medium <i>F. Bassani, M. Artoni, G.C. La Rocca, and I. Carusotto</i>	85
Coupled plasmon-LO phonon modes at high magnetic fields <i>A. Wyszynolek, M. Potemski, and T. Shupinski</i>	87
Excitation induced flips of the phase of transmitted pulses <i>G. Manzke, K. Henneberger, J.S. Nägerl, B. Stabenau, G. Böhne, and R.G. Ulbrich</i>	89
Below bottleneck polaritonic radiation in ultra high quality AlGaAs alloys <i>T.S. Shamirzaev, A.I. Toropov, A.K. Bakarov, K.S. Zhuravlev, A.Yu. Kobitsky, H.P. Wagner, and D.R.T. Zahn</i>	91
Coherent control of speckle: Resonant Rayleigh scattering from a conjugated polymer <i>S.P. Kennedy, N. Garro, and R.T. Phillips</i>	93
Fine structure of excitons in diamond: A cathodoluminescence study <i>N. Teofilov, R. Sauer, K. Thonke, T.R. Anthony, and H. Kanda</i>	95
Transient reflection from anisotropic semiconductors studied with coherent control method <i>Y. Mitsumori, M. Sato, H. Tobioka, and F. Minami</i>	97
Exciton-phonon droplets with Bose-Einstein condensate: transport and optical properties <i>D. Roubtsov, Y. Lépine, and I. Loutsenko</i>	99
Role of the excitonic continuum in the polariton problem <i>E.A. Muljarov and R. Zimmermann</i>	101
Far IR radiation from p-Ge under crossed directions of electric field and uniaxial pressure <i>A.A. Abramov, V.I. Akimov, A.T. Dalakyan, V.N. Tulupenko, D.A. Firsov, V.M. Bondar, V.N. Poroshin, V.I. Gavrilenko, and M.S. Kagan</i>	103
Anomalous transport of excitons in Cu ₂ O <i>S.G. Tikhodeev and N.A. Gippius</i>	105
Pump-probe spectroscopy by infrared probe light in highly excited semiconductors: Beyond the BCS-like mean-field theory <i>T.J. Inagaki and M. Aihara</i>	107
Strain-and field-induced optical anisotropies of GaAs measured by RDS <i>N. Kumagai, T. Yasuda, T. Hanada, and T. Yao</i>	109
Electromagnetically induced transparency for intrinsic exciton lines in bulk semiconductors <i>M. Artoni, G.C. La Rocca and F. Bassani</i>	111
Reversible photoinduced structural changes in GeSe ₂ glass at low-temperature <i>T. Nakaoka, Y. Wang, O. Matsuda, K. Inoue, and K. Murase</i>	113
Photoluminescence due to the scattering process from heavy-hole to light-hole excitons in CuI thin films <i>I. Tanaka and M. Nakayama</i>	115
Identification of fine structures of magneto-oscillatory spectra in cuprous oxide as quantum manifestation of classical non-integrability <i>K. Hammura, K. Sakai, and M. Seyama</i>	117
Optical studies of alloy semiconductors Ge _{1-x} C _x ($x < 0.05$) grown on Si substrates by combined low-energy ion beam and molecular beam epitaxy <i>M. Sakai, Y. Kitayama, S. Takaku, S. Kimura, and H. Shibata</i>	119

Optical properties of Ga ₂ Se ₃ <i>H. Nishimura, M. Takumi, K. Nagata, and Y. Miyamoto</i>	121
Exciton sublattice of incommensurate phase in low-dimensional TlInS ₂ <i>N. Mamedov, T. Aoki-Matsumoto, B. Gadjiev, H. Uchiki, N. Yamamoto, and S. Iida</i>	123
2.4 Transport and Localization	
Isotopic mass and lattice constants of Si and Ge: X-ray standing wave measurements (invited) <i>J. Zegenhagen, A. Kazimirov, L.X. Cao, M. Konuma, E. Sozontov, D. Plachke, H.D. Carstanjen, G. Bilger, E. Haller, V. Kohn, and M. Cardona</i>	125
Metal-insulator transition in doped semiconductors (invited) <i>K.M. Itoh</i>	128
Isotope effect on the thermal conductivity of silicon <i>T. Ruf, R.W. Henn, M. Asen-Palmer, E. Gmelin, M. Cardona, H.-J. Pohl, G.G. Devyatych, P.G. Sennikov, and J. Bollmann</i>	132
Axisymmetric Gunn effect <i>L.L. Bonilla, R. Escobedo, and F.J. Higuera</i>	134
DC field response of hot carriers under circular polarized intense microwave fields in semiconductors <i>N. Ishida</i>	136
Transient photoconductivity of pure CuO <i>H. Yamaguchi, T. Ito, and T. Masumi</i>	138
A microscopic simulation of the evolution of local perturbations <i>P. Gaubert, L. Varani, J.C. Vaissière, J.P. Nougier, E. Starikov, P. Shiktorov, and V. Gruzinskis</i>	140
Dispersive energy transport, relaxation current and Einstein relationship in band tails <i>O. Blcibaum, H. Böttger, V.V. Bryksin, and A.N. Samukhin</i>	142
Angle-resolved tunneling spectroscopy of Si conduction band using bonded Si(111) wafer pair <i>A. Kazama, H. Bando, Y. Miyahara, H. Enomoto, and H. Ozaki</i>	144
Identification of physical mechanisms of tensoeffects in highly strained Si and Ge <i>S.I. Budzulyak, V.M. Ermakov, Yu.P. Dotsenko, V.V. Kolomoets, Yu.M. Shwarts, E.F. Venger, M. Fukuzawa, P. Verma, M. Yamada, E. Liarokapis, and D.P. Tunstall</i>	146
Metal-insulator transition in anisotropic systems <i>F. Milde, R.A. Römer, and M. Schreiber</i>	148
Hot electron degradation of the extraordinary magneto-resistance in inhomogenous InSb <i>D. Poplavskyy, A.C.H. Rowe, R.A. Stradling, and S.A. Solin</i>	150
Critical exponents for the metal-insulator transition of ⁷⁰ Ge:Ga in magnetic fields <i>M. Watanabe, K.M. Itoh, M. Morishita, Y. Ootuka, and E.E. Haller</i>	152
Boron in diamond at high concentrations <i>M.J.R. Hoch, J.E. Lowther, and T. Tshepe</i>	154
Impurity scattering in metallic carbon nanotubes with superconducting pairs <i>K. Harigaya</i>	156
Self-organization of current density filaments in <i>n</i> -GaAs Corbino discs <i>K. Aoki</i>	158
Electrical observation of impurity levels in boron doped homoepitaxial diamond <i>T. Inushima, R.F. Mamin, T. Matsushita, S. Ohoya, and H. Shiomi</i>	160
Monte Carlo simulation of temperature-and field-dependent impact ionization for GaAs <i>H.K. Jung, S.W. Ko, C.K. You, and K. Taniguchi</i>	162
A study on temperature-and field-dependent impact ionization coefficient for silicon using Monte Carlo simulation <i>H.K. Jung, C.K. You, S.W. Ko, and K. Taniguchi</i>	164

Critical behavior of the thermoelectric transport properties in amorphous systems near the metal-insulator transition <i>C. Villagonzalo, R.A. Römer, M. Schreiber, and A. MacKinnon</i>	166
Photo-quenching in small-polaronic conduction in LiMn_2O_4 <i>K. Kushida and K. Kuriyama</i>	168
Low-frequency oscillations and chaos in semi-insulating GaAs subjected to a high magnetic field <i>A. Neumann, A.G.M. Jansen, P. Wyder, and R. Deltour</i>	170
The influence of the temperature in the scattering mechanisms in pure GaSb <i>L.G.O. Messias and E. Marega Jr.</i>	172
2.5 Carrier Relaxation and Ultrafast Phenomena	
Reversible quantum dynamics of impurity-bound electrons in GaAs (invited) <i>B.E. Cole, J.B. Williams, M.S. Sherwin, and C.R. Stanley</i>	174
Magnetic field enhanced terahertz emission from semiconductor surfaces <i>R. McLaughlin, A. Corchia, M.B. Johnston, C.M. Ciesla, D.D. Arnone, G.A.C. Jones, E.H. Linfield, A.G. Davies, and M. Pepper</i>	178
Dephasing of coherent LO phonon-plasmon coupled modes in polar semiconductors <i>S. Katayama, M. Hase, M. Iida, and S. Nakashima</i>	180
Femtosecond spectroscopy of large-momentum excitons in GaAs <i>M. Betz, G. Göger, A. Leitenstorfer, M. Bichler, W. Wegscheider, and G. Abstreiter</i>	182
A novel biexcitonic, non-radiative electron-hole recombination mechanism and its application in hydrogenated silicon semiconductors <i>S.B. Zhang and H.M. Branz</i>	184
Annihilation of coherent LO phonon-plasmon coupled modes by lattice defects in n-GaAs <i>M. Hase, K. Ishioka, K. Ushida, and M. Kitajima</i>	186
Ultrafast electro-absorption at the transition between classical and quantum response <i>J. Kono and A.H. Chin</i>	188
Deviations from diffusive transport in 100 nm - 100 fs spatio-temporal pump-probe experiments on GaAs <i>J. Hetzler, A. Brunner, M. Wegener, S. Leu, S. Nau, and W. Stolz</i>	190
Femtosecond optical excitation of acoustic phonons in bulk GaAs <i>O.B. Wright, B. Perrin, O. Matsuda, and V.E. Gusev</i>	192
Theory of hot luminescence in pulse-excited semiconductors <i>K. Hannewald, S. Glutsch, and F. Bechstedt</i>	194
Dynamics of Bloch oscillations under the influence of scattering <i>F. Löser, Yu.A. Kosevich, K. Köhler, and K. Leo</i>	196
Nonlinear response due to propagating excitonic polaritons in GaAs heterostructure <i>K. Akiyama, N. Tomita, Y. Nishimura, Y. Nomura, and T. Isu</i>	198
Dynamic inter-sideband Fano interference of excitons in ac-driven superlattices <i>R.-B. Liu and B.-F. Zhu</i>	200
Ultrafast inter-conduction band carrier dynamics in SiC <i>T. Tomita, S. Saito, T. Suemoto, H. Harima, and S. Nakashima</i>	202
Time-resolved stimulated photon echoes in layered semiconductors <i>H. Tobioka, Y. Mitsumori, and F. Minami</i>	204
Two-color time-resolved electronic Raman measurement in semiconductors <i>S. Saito and T. Suemoto</i>	206
Fermi edge singularities in out-of-equilibrium semiconductors <i>M. Combescot, B. Roulet, and C. Tanguy</i>	208

Theory of coherent oscillation of phonon-polaritons <i>C.S. Kim, A.M. Satanin, G.D. Sanders, and C.J. Stanton</i>	210
2.6 Phase Transition and Ordering	
Self-organized electron density patterns in n-GaAs induced by microwaves <i>V. Novák, V.V. Bel'kov, D.M. Mathina, S.D. Ganichev, and W. Prettl</i>	212
Novel observation of magnetic-field induced metal-insulator transition in n-GaAs through far-infrared photoconductivity measurements <i>H. Kobori, M. Inoue, and T. Ohyama</i>	214
Magnetization of single MnTe (sub)monolayers embedded in nonmagnetic quantum wells <i>G. Prechtl, W. Heiss, S. Mackowski, A. Bonanni, E. Janik, H. Sitter, and W. Jantsch</i>	216
Energy relaxation by phonon scattering in long-range ordered $(\text{Al}_{0.5}\text{Ga}_{0.5})_{0.5}\text{In}_{0.5}\text{P}$ <i>T. Kita, K. Yamashita, T. Nishino, Y. Wang, and K. Murase</i>	218
Low-dimensional magnetism in polymorphic compounds, VOMoO_4 and $(\text{V}_{0.56}, \text{Mo}_{0.44})_2\text{O}_5$ <i>I. Shiozaki, S. Takada, S. Wada, and T. Toyoda</i>	220
Highly efficient luminescence in partially ordered GaInP_2 <i>X.H. Zhang, S.J. Chua, and J.R. Dong</i>	222
Kinetics of the amorphous-to-crystalline phase transformation in Ge:Sb:Te alloys used for the phase-change optical memory applications <i>E. García-García, E. Morales-Sánchez, J. González-Hernández, E. Prokhorov, Yu. Vorobiev, and S. Kostylev</i>	224
The intensity and wavelength dependence for photo-induced crystallization process in amorphous GeSe_2 <i>K. Sakai, K. Maeda, H. Yokoyama, and T. Ikari</i>	226
Stiffness transition and connectivity in amorphous chalcogenide semiconductors <i>Y. Wang, T. Nakaoka, M. Nakamura, O. Matsuda, and K. Murase</i>	228
Effect of pressure on electrical transport in electron doped perovskite manganite $\text{Sr}_{0.9}\text{Ce}_{0.1}\text{MnO}_3$ <i>T. Eto, F. Honda, G. Oomi, and A. Sundaresan</i>	230
2.7 Magnetic and Semimagnetic Semiconductors	
Ferromagnetism in diluted magnetic semiconductors <i>J. König, H.-H. Lin, and A.H. MacDonald</i>	232
Ferromagnetic interactions in p- and n-type II-VI diluted magnetic semiconductors <i>T. Andrearczyk, J. Jaroszyński, M. Sawicki, L. van Khoi, T. Dietl, D. Ferrand, C. Bourgognon, J. Cibert, S. Tatarenko, T. Fukumura, Z. Jin, H. Koinuma, and M. Kawasaki</i>	234
Theoretical predictions for transparent ferromagnets with transition metal atom doped ZnO <i>K. Sato and H. Katayama-Yoshida</i>	236
<i>sp-d</i> exchange interaction in GaMnAs investigated by resonant Kerr effect under high magnetic field <i>F. Michelini, N. Nègre, G. Fishman, M. Goiran, J. Sadowski, E. Vanelle, and S. Askénasy</i>	238
Giant tunability of excitonic photoluminescence transitions in antiferromagnetic EuTe epilayers <i>W. Heiss, G. Prechtl, and G. Springholz</i>	240
Hybridization-induced exchange interaction between the conduction band electrons and Mn ions in diluted magnetic semiconductors <i>A.K. Bhattacharjee and J. Pérez-Conde</i>	242
Cross-sectional scanning tunneling microscope (STM) study of Mn-doped GaAs layers <i>T. Tsuruoka, R. Tanimoto, N. Tachikawa, S. Ushioda, F. Matsukura, and H. Ohno</i>	244
Anomalous enhancement of the cyclotron mass of electrons in $\text{Cd}_{1-x}\text{Mn}_x\text{Te}$ observed at very high magnetic fields <i>Y.H. Matsuda, T. Ikaida, T. Yasuhira, N. Miura, S. Kuroda, F. Takano, and K. Takita</i>	246

Electronic structure of superlattices of II–VI/III–V diluted magnetic semiconductors <i>T. Kamatani and H. Akai</i>	248
Tunneling into amorphous Gd_xSi_{1-x} at the metal-insulator transition and its independence of magnetic impurities in the barrier <i>W. Teizer, F. Hellman, and R.C. Dynes</i>	250
Spin-lattice relaxation in semimagnetic quantum wells with a 2DEG <i>D.R. Yakovlev, A.V. Scherbakov, B. König, W. Ossau, A.V. Akimov, T. Wojtowicz, G. Karczewski, and J. Kossut</i>	252
Magnetic-field-driven metal-insulator transition in magnetic semiconductor (Ga,Mn)As <i>T. Hayashi, Y. Hashimoto, S. Yoshida, S. Katsumoto, and Y. Iye</i>	254
The effect of stress on magnetic properties of ferromagnetic EuS-PbS and EuS-PbSe semiconductor structures <i>T. Story, M. Arciszewska, M. Chernyshova, W. Dobrowolski, W. Mac, A. Twardowski, R. Świrkwicz, and A. Yu. Sipatov</i>	256
Fundamental properties of Fe-based III-V magnetic alloy semiconductor (Ga, Fe)As <i>R. Moriya, T. Kondo, Y. Katsumata, H. Munekata and S. Haneda</i>	258
The waves of spin and charge densities in diluted magnetic semiconductors near the paramagnetic-ferromagnetic phase transition <i>Yu.G. Semenov and V.N. Sokolov</i>	260
Electronic structure of $Ga_{1-x}Mn_xAs$ studied by photoemission spectroscopy <i>J. Okabayashi, A. Kimura, O. Rader, T. Mizokawa, A. Fujimori, T. Hayashi, and M. Tanaka</i>	262
Electronic structure and magnetic properties of (Zn,Mn)Se with carriers doping studied by FLAPW method <i>K. Matsushita, H. Harima, A. Yanase, and H. Katayama-Yoshida</i>	264
Cyclotron resonance of spin polaron in CdMnTe/CdMgTe 2D electron systems <i>Y. Imanaka, T. Takamasu, G. Kido, G. Karczewski, T. Wojtowicz, and J. Kossut</i>	266
Far infrared absorption spectra in ferromagnetic $Ga_{1-x}Mn_xAs$ <i>Y. Nagai, T. Kunimoto, K. Nagasaka, H. Nojiri, M. Motokawa, F. Matsukura, and H. Ohno</i>	268
Dynamical aspects of exciton magnetic polaron formation in diluted magnetic semiconductor quantum wells <i>T. Stirner and W.E. Hagston</i>	270
Cryobaric exciton magnetophotoluminescence in $Cd_{1-x}Mn_xSe$ <i>N. Kuroda and Y.H. Matsuda</i>	272
Line shape of excitation spectra of photoluminescence in CdMnTe and its quantum wells <i>K. Takamura, S. Yamamoto, and J. Nakahara</i>	274

3. Growth, Surfaces, and Interfaces

3.1 Atomic Structure

Possibility of the quantum fluctuation of the Si(001) surface at low temperature <i>Y. Yoshimoto and M. Tsukada</i>	277
(2×4) and (4×2) reconstructions of GaAs(001): The surface phase diagram re-examined <i>W.G. Schmidt, S. Mirbt, and F. Bechstedt</i>	279
Theory of $Al_2O_3(0001)$ surfaces and their employment as a substrate for nitride growth <i>R. Di Felice and J.E. Northrup</i>	281
Influence of strain and diffusion on the growth of V-groove InGaAs/GaAs quantum wires <i>K. Leifer, F. Lelarge, A. Rudra, S. Stauss, and E. Kapon</i>	283
Atomic structure at a Si (001) oxidation front <i>N. Ikarashi, K. Watanabe, and Y. Miyamoto</i>	285

Binding and migration paths of Au adatoms on the GaAs (001)- $\beta 2(2 \times 4)$ surface <i>A.A. Bonapasta and F. Buda</i>	287
Surface topography of Si(111)- 7×7 reconstruction: first-principles investigations <i>S.H. Ke, T. Uda, and K. Terakura</i>	289
A structural model of Si(111)($2\sqrt{3} \times 2\sqrt{3}$) R30°-Sn <i>T. Ichikawa, K. Cho, T. Onodera, A. Mizoguchi, and T. Ohkoshi</i>	291
Synchrotron radiation photoemission studies of surface reconstruction on GaAs (001) <i>K. Ono, T. Mano, K. Nakamura, M. Mizuguchi, S. Nakazono, K. Horiba, T. Kihara, H. Kiwata, I. Waki, M. Oshima, N. Koguchi, and A. Kakizaki</i>	293
Structural ordering on Si(111) $\sqrt{3} \times \sqrt{3}$ -Ag surface : Monte Carlo simulation based on first-principles calculations <i>Y. Nakamura, Y. Kondo, J. Nakamura, and S. Watanabe</i>	295
STM and RHEED study of Ge(110) reconstructions <i>T. Ichikawa, H. Fujii, and A. Sugimoto</i>	297
3.2 Electronic Structure	
Terrace and step contributions to the surface optical anisotropy of Si(001) <i>W.G. Schmidt, F. Bechstedt, and J. Bernholc</i>	299
Nano-scale ferromagnets on semiconductors: Ga adsorbates on Si (100) surfaces <i>S. Okada and A. Oshiyama</i>	301
AlGaIn/GaN lateral polarity heterostructures <i>A.P. Lima, C. Miskys, O. Ambacher, M. Stutzmann, R. Dimitrov, V. Tilak, M.J. Murphy, and L.F. Eastman</i>	303
Modification of the Si surface electronic properties by Ge nanostructures: Surface photovoltage studies <i>K. Nauka and T.I. Kamins</i>	305
<i>Ab initio</i> study of SiC/Ti polar interfaces <i>S. Tanaka and M. Kohyama</i>	307
3.3 Adsorption and Surface Reactions	
Microscopic mechanism of Si oxidation (invited) <i>K. Shiraishi, H. Kageshima, and M. Uematsu</i>	309
Electron paramagnetic resonance of a single-crystal surface: the Si(111)- 7×7 surface and its oxidation process <i>T. Umeda, M. Nishizawa, T. Yasuda, J. Isoya, S. Yamasaki, and K. Tanaka</i>	313
Atomic layer oxidation of H terminated Si(100) surface : Domino reaction via oxidation and H migration <i>K. Kato, H. Kajiyama, S. Heike, T. Hashizume, and T. Uda</i>	315
Time-dependent density-functional simulations of desorption dynamics of H and Br terminated Si surfaces induced by electronic excitations <i>Y. Miyamoto and O. Sugino</i>	317
Mechanism of migration and dissociation for As ₂ molecule on GaAs(001) surfaces during the MBE growth: A computational study <i>A. Ishii, K. Seino, and T. Aisaka</i>	319
Photoemission spectroscopy of Si(001) surfaces oxidized by hyperthermal O ₂ molecular beams <i>Y. Teraoka and A. Yoshigoe</i>	321
Formation of first InSb molecular layer on Si(111) substrate: Role of In(4 \times 1) reconstruction <i>B.V. Rao, D. Gruznev, T. Tambo, and C. Tatsuyama</i>	323
Laser-induced electronic instability on semiconductor surfaces of Si (111)-(7 \times 7) and InP (110)-(1 \times 1) <i>K. Tanimura, J. Kanasaki, and K. Ishikawa</i>	325

Surface states of (100)n-GaAs with adsorbed oxygens and their dependence on chemical treatment <i>Y. Kasai, T. Tsuzuku, Y. Ohta, T. Inokuma, K. Iiyama, and S. Takamiya</i>	327
Real-time observation of initial oxidation on highly B-doped Si (100)-2×1 surfaces using scanning tunneling microscopy <i>K. Ohmori, M. Tsukakoshi, H. Ikeda, A. Sakai, S. Zaima, and Y. Yasuda</i>	329
Fabrication of boron delta-doped structures in Si by solid phase epitaxy <i>T. Ishikawa, H. Nagai, K. Ishii, and S. Matsumoto</i>	331
Effect of gas adsorption on the interfacial states at n-SnO ₂ /p-Si heterojunction <i>R.B. Vasiliev, A.M. Gaskov, M.N. Rumyanseva, and L.I. Ryabova</i>	333
Structure of a SiN layer on Si(111) surface <i>T. Yamasaki, C. Kaneta, and T. Uda</i>	335
3.4 Bulk Growth and Thin Film Epitaxy	
Distinct morphological evolution of Si _{1-x} Ge _x films on Si(100) during gas-source MBE and photo-CVD <i>H. Akazawa</i>	337
Toward predictive growth simulations: MBE on GaAs(001) <i>P. Kratzer and M. Scheffler</i>	339
Dual Au-Si bonding character and Au surfactant effect on Au/Si(111) surfaces <i>M. Murayama, T. Nakayama, and A. Natori</i>	341
Theoretical study on epitaxial growth of lattice-mismatched semiconductor systems <i>N. Miyagishima, K. Okajima, K. Takeda, N. Oyama, T. Ohno, K. Shiraishi, and T. Ito</i>	343
Control of electronic structure and thermodynamic stability of SrTiO ₃ with Ru or Nb doping from first-principles calculation <i>T. Shimizu and T. Kawakubo</i>	345
Modeling and simulation of FBAR devices fabricated on Si <i>D.H. Kim, G. Yoon, and H.D. Park</i>	347
Nucleation mechanism of stacking fault during Si epitaxial growth by chemical vapor deposition with dichlorosilane <i>Y. Takakuwa, M.K. Mazumder, and N. Miyamoto</i>	349
Self-limiting etching prior to self-limiting growth in ultra-high vacuum for obtaining clean interface <i>N. Otsuka, J. Nishizawa, Y. Oyama, H. Kikuchi, and K. Suto</i>	351
Reflectance-difference spectroscopy of (001) InAs surfaces in ultrahigh vacuum <i>T. Kita, H. Tango, K. Tachikawa, K. Yamashita, and T. Nishino</i>	353
3.5 Growth of Self-Assembled Dots	
Controlling of lateral and vertical order in self-organized PbSe quantum dot superlattices (invited) <i>G. Springholz, M. Pinczolits, V. Holy, P. Mayer, G. Bauer, H. H. Kang, and L. Salamanca-Riba</i>	355
Shape analysis of single and stacked InAs quantum dots at the atomic level by cross-sectional STM <i>D.M. Bruls, J.W.A.M. Vugs, P.M. Koenraad, M.S. Skolnick, M. Hopkinson, and J.H. Wolter</i>	359
Formation process of InAs dots including Mn atoms and their physical properties <i>S. Okumura, H. Asahi, Y.K. Zhou, J. Asakura, M. Kanamura, K. Asami, H. Kubo, C. Hamaguchi, and S. Gonda</i>	361
Shape and size of buried SiGe islands <i>J. Stangl, V. Holý, A. Daniel, T. Roch, G. Bauer, T.H. Metzger, J. Zhu, K. Brunner, and G. Abstreiter</i>	363
Optical anisotropy of Stranski-Krastanov growth surface of InAs on GaAs (001) <i>T. Kita, H. Tango, K. Tachikawa, K. Yamashita, T. Nishino, T. Nakayama, and M. Murayama</i>	365
Lasers based on self-assembled InAs/GaAs and InP/InGaP quantum dots <i>O.G. Schmidt, M.O. Lipinski, Y.M. Manz, H. Heidemeyer, W. Winter, and K. Eberl</i>	367

Effect of vertical size uniformity on diffraction contrast images of stacked InGaAs/GaAs quantum dots <i>M. De Giorgi, A. Passaseo, R. Cingolani, A. Taurino, and M. Catalano</i>	369
Lateral distribution of buried self-assembled InAs quantum dots on GaAs <i>K. Zhang, J. Falta, Ch. Heyn, Th. Schmidt, and W. Hansen</i>	371
Nucleation site control in self-assembling Si quantum dots on ultrathin SiO ₂ /c-Si <i>S. Miyazaki, M. Ikeda, E. Yoshida, N. Shimizu, and M. Hirose</i>	373
Reversibility of the island shape, volume and density in Stranski-Krastanow growth <i>N.N. Ledentsov, V.A. Shchukin, R. Heitz, D. Bimberg, V.M. Ustinov, N.A. Cherkashin, A.R. Kovsh, Yu.G. Musikhin, B.V. Volovik, A.E. Zhukov, G.E. Cirlin, and Zh.I. Alferov</i>	375
Effect of Si diffusion on growth of GeSi self-assembled islands <i>A.V. Novikov, N.V. Vostokov, S.A. Gusev, Yu.N. Drozdov, Z.F. Krasil'nik, D.N. Lobanov, L.D. Moldavskaya, M. Miura, V.V. Postnikov, M.V. Stepikhova, Y. Shiraki, and N. Usami</i>	377
Observation of a universal behavior in the growth of InAs self-assembled quantum dots on patterned substrates <i>S.W. Hwang, M.H. Son, B.H. Choi, D. Ahn, C.K. Hyon, S.-H. Song, Y.J. Park, and E.K. Kim</i>	379
Monte Carlo simulation of the self-organised growth of quantum dots with anisotropic surface diffusion <i>M. Meixner, R. Kunert, S. Bose, E. Schöll, V.A. Shchukin, D. Bimberg, E. Penev, and P. Kratzer</i>	381
Growth of InAs quantum dots on (110) -oriented cleaved GaAs surfaces <i>M. Gerling, S. Jeppesen, A. Gustafsson, and L. Samuelson</i>	383
Fabrication of compound-semiconductor quantum dots on a Si(111) substrate terminated by bilayer-GaSe <i>K. Ueno, K. Saiki, and A. Koma</i>	385
Optimized growth procedure for self-organized InAs quantum dots <i>E. Steimetz, T. Wehnert, P. Kratzer, L.G. Wang, Q.K.K. Liu, H. Kirmse, J.-T. Zettler, W. Neumann, M. Scheffler, and W. Richter</i>	387
Effect of inter-island interaction on the growth of self-assembled quantum dots <i>S. Vannarat, S.T. Chui, K. Esfarjani, and Y. Kawazoe</i>	389
Properties of CdSe/ZnSe based quantum heterostructures with and without lateral confinement potentials <i>E. Kurtz, M. Schmidt, B. Dal Don, S. Wachter, D. Litvinov, D. Gerthsen, H. Kalt, and C. Klingshirn</i>	391
Growth, morphology and optical properties of metal clusters on semiconductor surface (Au/GaAs) <i>N. Dmítruk, T. Barlas, I. Dmitruk, T. Mikhailik, and V. Romaniuk</i>	393
Formation of GaAsN nanoinclusions in a GaN matrix <i>A.F. Tsatsul'nikov, I.L. Krestnikov, W.V. Lundin, A.V. Sakharov, D.A. Bedarev, A.S. Usikov, B. Ya. Ber, V.V. Tret'yakov, Zh.I. Alferov, N.N. Ledentsov, A. Hoffmann, D. Bimberg, T. Riemann, J. Christen, Yu.G. Musikhin, I.P. Soshnikov, D. Litvinov, A. Rosenauer, D. Gerthsen, and A. Plaut</i>	395
Extremely uniform InAs/GaAs quantum dots emitting at 1.46 μm at room temperature grown by MOCVD with Bi doping <i>B.N. Zvonkov, I.A. Karpovich, N.V. Baidus, D.O. Filatov, Yu.Yu. Gushina, S.V. Morozov, and S.B. Levichev</i>	397
Effects of InAs coverage on the Ga diffusion into InAs self-assembled quantum dots on GaAs(100) <i>N. Matsumura, T. Haga, S. Muto, Y. Nakata, and N. Yokoyama</i>	399
Photoluminescence and atomic force microscopy studies of InAs/InSb nanostructures grown by MBE <i>Ya.V. Terent'ev, A.A. Toropov, V.A. Solov'ev, B. Ya. Mel'tser, M.M. Moiseeva, S.V. Ivanov, B. Magnusson, B. Monemar, and P.S. Kop'ev</i>	401
Three dimensional self-organization of InAs quantum-dot multilayers <i>J.C. González, F.M. Matinaga, W.N. Rodrigues, M.V.B. Moreira, A.G. de Oliveira, M.I.N. da Silva, J.M.C. Vilela, M.S. Andrade, D. Ugarte, and P.C. Silva</i>	403
Investigation of the properties of molecular beam epitaxy grown self-organised ZnSe quantum dots embedded in ZnS <i>C.L. Yang, L.W. Lu, W.K. Ge, Z.H. Ma, I.K. Sou, and J.N. Wang</i>	405

Exciton localization in alloy/alloy interfaces of InGaAs/GaAs(001) stepped quantum wells <i>A. D'Andrea, F. Fernández-Alonso, M. Righini, D. Schiumarini, S. Selci, and N. Tomassini</i>	407
Characterization of Ge nanocrystal embedded in SiO ₂ films <i>H. Fukuda, S. Sakuma, T. Yamada, S. Nomura, M. Nishino, T. Higuchi, and S. Ohshima</i>	409
3.6 Semiconductor–Nonsemiconductor Interfaces	
Control of quantum confinement in metal-clad InAs quantum wells <i>S. Tsujino, S.J. Allen, M. Rügenacht, M. Thomas, J.P. Zhang, J. Speck, T. Eckhause, and B. Gwinn</i>	411
Unsaturated cyclic-hydrocarbon molecules on a Si (001) surface: A first-principles approach <i>K. Akagi and S. Tsuneyuki</i>	413
Organic modified GaAs(100) Schottky contacts <i>Th. Lindner, T.U. Kampen, S. Park, and D.R.T. Zahn</i>	415
Interface effects on exciton states in a CdTe/(Cd, Mn)Te quantum well <i>H. Yokoi, Y. Kakudate, Yu.G. Semenov, S. Takeyama, S.W. Tozer, Y. Kim, T. Wojtowicz, G. Kar-czewski, and J. Kossut</i>	417
Reaction of atomic hydrogen with the Si(100)/SiO ₂ interface defects <i>C. Kaneta, T. Yamasaki, and T. Uda</i>	419
Temperature dependence of low frequency noise mechanisms in Schottky barrier structure <i>J.I. Lee, I.K. Han, J. Brini, A. Chovet, and C.A. Dimitriadis</i>	421
Interfacial transition regions of gate dielectrics in advanced silicon devices <i>G. Lucovsky, J.C. Phillips, and M.F. Thorpe</i>	423
ESR and magnetic measurement of Ni/GaAs composite system <i>Y. Seino, S.A. Haque, A. Matsuo, Y. Yamamoto, S. Yamada, and H. Hori</i>	425
3.7 STM, AFM, and Near-Field Studies of Surfaces and Interfaces	
Imaging of Friedel oscillations at epitaxially grown InAs(111)A surfaces using scanning tunneling microscopy (invited) <i>K. Kanisawa, M.J. Butcher, H. Yamaguchi, and Y. Hirayama</i>	427
Probing of subsurface dopants buried in silicon by scanning tunneling microscopy <i>Y. Suwa, T. Hitosugi, S. Matsuura, S. Heike, S. Watanabe, T. Onogi, and T. Hashizume</i>	431
Effect of tip morphology on AFM images: <i>Ab initio</i> simulations on GaAs(110) surface <i>S.H. Ke, T. Uda, I. Štich, and K. Terakura</i>	433
Novel investigation technique for interior III/V-semiconductor interfaces <i>S. Nau, G. Bernatz, and W. Stolz</i>	435
Observation of dopant-atom dimers on hydrogen-terminated Si(100)-2×1 surface by scanning tunneling microscopy <i>S. Matsuura, M. Fujimori, S. Heike, Y. Suwa, T. Onogi, H. Kajiyama, K. Kitazawa, and T. Hashizume</i>	437
Reconstruction on Si(100) surface induced by the type-A defects near T_c <i>M. Okamoto, T. Yokoyama, and K. Takayanagi</i>	439
Space and energy distribution of surface gap states on MBE-grown and silicon-covered (001) GaAs surfaces studied by scanning tunneling spectroscopy <i>N. Negoro, S. Kasai, and H. Hasegawa</i>	441
Atomically resolved imaging of semiconductor surfaces using noncontact atomic force microscopy <i>S. Morita and Y. Sugawara</i>	443
Atomic structure of GaP (001) and InP (001) reconstructions: Scanning tunneling microscopy and <i>ab initio</i> theory <i>K. Lüdge, P. Vogt, O. Pulci, N. Esser, F. Bechstedt, and W. Richter</i>	445

Bias voltage dependence of scanning tunneling microscopy images of a Si (100) ₂ ×3-Ba surface <i>K. Ojima, M. Yoshimura, and K. Ueda</i>	447
Removal of particles from surface of silicon using clean solutions with surfactants and chelates <i>C. Baocheng, Y. Xinhao, M. Honglei, M. Jin, and L. Zhengli</i>	449
4. Heterostructures and Superlattices: Optical	
4.1 Structural Properties	
Phase difference between coherent GaSb-like and AlSb-like LO phonons in GaSb/AlSb superlattices <i>H. Takeuchi, K. Mizoguchi, M. Nakayama, K. Kuroyanagi, and T. Aida</i>	451
Plasmon-phonon coupling at Ga _{0.5} In _{0.5} P/GaAs heterointerfaces induced by CuPt-type ordering <i>K. Yamashita, T. Kita, T. Nishino, Y. Wang, K. Murase, C. Geng, F. Scholz, and H. Schweizer</i>	453
Optical properties and band alignment of ZnSe/ZnMgBeSe heterostructures <i>K. Godo, M.W. Cho, J.H. Chang, S.K. Hong, H. Makino, T. Yao, M.Y. Shen, and T. Goto</i>	455
Analysis of Raman spectra of quasiperiodic GaAs-AlAs heterostructures <i>E. Matsushita and T. Furuyama</i>	457
4.2 Electronic Structures	
Controlling Fermi-edge singularities by a periodic external potential <i>S. Nomura, T. Nakanishi, and Y. Aoyagi</i>	459
Optical anisotropy change of buried SiO ₂ /Si interfaces during layer-by-layer oxidation <i>T. Nakayama and M. Murayama</i>	461
Negatively charged excitons in semiconductor quantum wells: effects of longitudinal electric and magnetic fields <i>L.C.O. Dacal, M.J.S.P. Brasil, and J.A. Brum</i>	463
The influence of intraband transitions on the resonant and extended electronic states in GaAs/AlGaAs asymmetric quantum wells as a function of their confinement <i>M. Levy, R. Beserman, R. Kapon, A. Sa'ar, V. Thierry-Mieg, and R. Planel</i>	465
Zeeman mapping of probability densities in square quantum wells using ultranarrow probes <i>G. Prechtl, W. Heiss, A. Bonanni, E. Janik, S. Mackowski, G. Karczewski, and W. Jantsch</i>	467
Tight-binding description of GaAs/AlAs quantum wells and superlattices: Gap transitions and intervalley couplings <i>R.B. Capaz, J.G. Menchero, T.G. Dargam, and B. Koiller</i>	469
Large quantum confinement effect of conduction electrons in ZnSe/BeTe type II heterostructures <i>R. Akimoto, Y. Kinpara, and K. Akita</i>	471
A study of the GaAs/partially ordered GaInP interface <i>T. Kobayashi, K. Inoue, A.D. Prins, K. Uchida, and J. Nakahara</i>	473
Analytical and transfer matrix solutions of the $\mathbf{k} \cdot \mathbf{p}$ Hamiltonian for the X-band in AlAs confined systems <i>L.E. Bremme and P.C. Klipstein</i>	475
Relation between the in-plane polarization anisotropy of the optical properties and the microscopic atomic configuration in (Ga _{0.5} In _{0.5} As)/(InP) superlattices <i>R. Magri and S. Ossicini</i>	477
Comparison of empirical pseudopotential and $\mathbf{k} \cdot \mathbf{p}$ calculations in p-doped strained layer SiGe quantum wells <i>Z. Ikonić, R.W. Kelsall, and P. Harrison</i>	479
Plasmons in laterally density modulated 2D electron gas in shallow etched single-heterostructures <i>T. Tagawa and S. Katayama</i>	481

Electronic properties of quantum wire superlattices elaborated by the 'Atomic Saw' method <i>F. Michelini, L. Ressler, G. Fishman, E. Vanelle, F. Laruelle, and J.P. Peyrade</i>	483
Cyclotron resonance of holes and shallow acceptor photoconductivity in strained MQW Ge/GeSi heterostructures in strong magnetic fields <i>V. Ya. Aleshkin, I.V. Erofeeva, V.I. Gavrilenko, O.A. Kuznetsov, M.D. Moldavskaya, V.L. Vaks, and D.B. Veksler</i>	485
Breakdown of rotational symmetry at semiconductor interfaces <i>O. Krebs, S. Cortez, and P. Voisin</i>	487
4.3 Magnetic Superlattices	
Binding energy and internal magnetic field of exciton magnetic polarons in a single semimagnetic quantum dot <i>G. Bacher, A.A. Maksimov, A. McDonald, M.K. Welsch, H. Schömiig, V.D. Kulakovskii, A. Forchel, Ch. Becker, G. Landwehr, and L.W. Molenkamp</i>	489
Spin transport of excitons in asymmetric double quantum wells of $\text{Cd}_{1-x}\text{Mn}_x\text{Te}$ <i>K. Kayanuma, E. Shirado, M.C. Debnath, I. Souma, S. Permogorov, and Y. Oka</i>	491
Raman scattering in MnTe/ZnTe and $\text{ZnSe}/\text{ZnMnSe}$ multilayer semiconductor superlattices <i>K. Ozawa, Y. Tanaka, K. Iio, N. Nakajima, Y. Nabetani, and T. Matsumoto</i>	493
4.4 Interband Transitions and Excitons	
Exciton-trion coupling in modulation doped quantum well structures <i>V.P. Kochereshko, G.V. Astakhov, R.A. Suris, D.R. Yakovlev, W. Ossau, J. Nurnberger, W. Faschinger, and G. Landwehr</i>	495
Rapid recombination process of free trions <i>D. Sanvitto, R.A. Hogg, A.J. Shields, D.M. Whittaker, M.Y. Simmons, D.A. Ritchie, and M. Pepper</i>	497
Spin dependent exciton-exciton interaction in hot and cold 2D exciton gases controlled by an electric field <i>G. Aichmayr, L. Viña, S.P. Kennedy, R.T. Phillips, and E.E. Mendez</i>	499
From excitons to Fermi edge singularity <i>G. Yusa, H. Shtrikman, and I. Bar-Joseph</i>	501
Magnetic traps for excitons in $\text{GaAs}/\text{Al}_x\text{Ga}_{1-x}\text{As}$ quantum wells <i>J.A.K. Freire, F.M. Peeters, A. Matulis, V.N. Freire, and G.A. Farias</i>	503
Excess carrier effects upon the excitonic absorption thresholds of remotely doped $\text{GaAs}/\text{AlGaAs}$ quantum wells <i>R. Kaur, A.J. Shields, R.A. Hogg, J.L. Osborne, M.Y. Simmons, D.A. Ritchie, and M. Pepper</i>	505
Observation of heavy-hole minibands in ultra-short period superlattices <i>M. Eckardt, W. Geisselbrecht, S. Malzer, G.H. Döhler, K. Maranowski, and A.C. Gossard</i>	507
CdTe quantum wells as ideal systems for the study of negatively charged excitons: A spectral and temporal analysis <i>V. Ciulin, P. Kossacki, M. Kutrowski, A. Esser, S. Haacke, J.-D. Gavière, T. Wojtowicz, and B. Deveaud</i>	509
Collapse of the excitonic states at $r_s=8$ in high quality $\text{GaAs}/\text{AlGaAs}$ single quantum wells <i>S.I. Gubarev, I.V. Kukushkin, S.V. Toustonog, M.Yu. Akimov, L.V. Kulik, J. Smet, K. von Klitzing, and W. Wegscheider</i>	511
Phase sensitive detection of transmitted femtosecond pulses in GaAs quantum wells <i>R. Kawahara, T. Kuroda, Y. Mitsumori, and F. Minami</i>	513
Oscillatory behavior of the Γ - X coupling with AlAs thickness in type II GaAs/AlAs heterostructures <i>C. Gourdon, D. Martins, V. Voliotis, P. Lavallard, and E.L. Ivchenko</i>	515
Blue Stark shift in composite quantum wells <i>A.Yu. Silov, B. Aneeshkumar, M.R. Leys, H. Vonk, and J.H. Wolter</i>	517

Spectroscopic studies of the interaction of an electron gas with photoexcited electron-hole pairs in modulation-doped GaAs/AlGaAs quantum wells <i>T. Yeo, H.A. Nickel, G. Comanescu, H.D. Cheong, M. Furis, A. Petrou, and B.D. McCombe</i>	519
Resonant Rayleigh scattering of exciton-polaritons in multiple quantum well structures <i>A.V. Kavokin, G. Malpuech, W. Langbein, and J.M. Hvam</i>	521
Speckle-correlation spectroscopy on localized spin-split exciton states in quantum wells <i>R. Zimmermann, W. Langbein, E. Runge, and J.M. Hvam</i>	523
Optical properties of trions in semiconductor nanostructures <i>A. Esser, E. Runge, and R. Zimmermann</i>	525
Internal transitions of charged magneto excitons in II-VI quantum well heterostructures <i>C.J. Meining, M. Furis, H.A. Nickel, D.R. Yakovlev, W. Ossau, A. Petrou, and B.D. McCombe</i>	527
Many-body corrections in n-type 2D telluride structures <i>V. Iuard, R.T. Cox, K. Saminadayar, C. Bourgognon, S. Tatarenko, and L. Besombes</i>	529
Possibility of excitonic polymers in type-II superlattices <i>T. Tsuchiya</i>	531
Localised exciton transitions in high-quality GaAs/AlGaAs quantum wells <i>A.G. Steffan, A. García-Cristóbal, R.T. Phillips, and D.A. Ritchie</i>	533
Field-induced optical anisotropy in semiconductor superlattices: the Wannier-Pockels effect <i>S. Cortez, O. Krebs, J.C. Harmand, J.L. Gentner, and P. Voisin</i>	535
Origin of optical-phonon Raman spectra in multiple quantum wells <i>B.-F. Zhu and S.F. Ren</i>	537
Temperature dependence of the band gap in (InGa)(AsN)/GaAs single quantum wells <i>A. Polimeni, M. Capizzi, M. Geddo, M. Fischer, M. Reinhardt, and A. Forchel</i>	539
Charge transfer of carriers by interband photoexcitation in asymmetric GaAs/AlGaAs coupled quantum wells <i>M. Levy, Yu.L. Khait, R. Beserman, A. Sa'ar, V. Thierry-Mieg, and R. Planel</i>	541
Optical properties of InGaAsN/GaAs QWs for long-wavelength lasers on GaAs <i>H. Riechert, A.Yu. Egorov, Gh. Dumitras, and B. Borchert</i>	543
Effective mass theory for a magneto-exciton in the type II superlattices or confinement structures and the origin of anomalous photoluminescence <i>K. Kouzu, M. Nishimura, and H. Kamimura</i>	545
Modulation spectroscopy of critical points in excitonic energy structure of thick GaAs quantum wells <i>V.G. Davydov, Yu.K. Dolgikh, Yu.P. Efimov, S.A. Eliseev, A.V. Fedorov, I.Ya. Gerlovin, I.V. Ignatiev, I.E. Kozin, V.V. Petrov, V.V. Ovsyankin, I.A. Yugova, H.-W. Ren, S. Sugou, and Y. Masumoto</i>	547
Exciton-phonon coupling in wide bandgap II-VI quantum wells <i>B. Urbaszek, A. Balocchi, C. Morhain, C. Bradford, X. Tang, C.M. Townsley, C.B. O'Donnell, S.A. Telfer, K.A. Prior, B.C. Cavenett, and R.J. Nicholas</i>	549
Magneto-optical evidence of many-body effects in spin-polarized 2D electron gas <i>C. Testelin, A. Lemaître, C. Rigaux, T. Wojtowicz, G. Karczewski, and F. Teran</i>	551
Forbidden interband transitions and the indirect valence band in strained GaInAs/InP quantum wells <i>A. Dörnen, J. Shao, V. Härle, and F. Scholz</i>	553
Magnetic field and pressure effects of photoluminescence in a GaP/AlP short-period superlattice <i>K. Uchida, N. Miura, F. Issiki, and Y. Shiraki</i>	555
Optical properties of interacting excitons in quantum wells <i>S. de-Leon and B. Laikhtman</i>	557
Recombination processes of GaNAs/GaAs structures : Effect of rapid thermal annealing <i>I.A. Buyanova, W.M. Chen, G. Pozina, P.N. Hai, N.Q. Thanh, H.P. Xin, and C.W. Tu</i>	559

Dressed excitons and shallow impurities in low-dimensional semiconductor systems within a renormalized effective-mass approach <i>H.S. Brandi, A. Latgé, and L.E. Oliveira</i>	561
Superlinear photoluminescence in GaAs/(GaAl)As heterojunctions <i>J.X. Shen, R. Pittini, and Y. Oka</i>	563
The exciton dead layer revisited <i>M. Combescot, R. Combescot, and B. Roulet</i>	565
Picosecond dynamics of polarized resonance photoluminescence in the GaAs/AlGaAs- superlattices <i>I.Ya. Gerlovin, Yu.K. Dolgikh, S.A. Eliseev, V.V. Ovsyankin, Yu.P. Efimov, I.V. Ignatiev, I.E. Kozin, V.V. Petrov, and Y. Masumoto</i>	567
Magneto-photorefectance of the above barrier state transitions in GaAs/Al _{0.3} Ga _{0.7} As double quantum wells <i>G. Sek, M. Nowaczyk, L. Bryja, K. Ryczko, J. Misiewicz, M. Bayer, J. Koeth, and A. Forchel</i>	569
Indirect transitions between barrier (X) electrons and two-dimensional hole gas in mixed type I - type II quantum wells <i>R. Guliamov, E. Lifshitz, E. Cohen, A. Ron, and L.N. Pfeiffer</i>	571
Electronic states of interface Al-2p core excitons in GaAs/AlAs/GaAs heterostructures <i>K. Inoue, Y. Ishiwata, and S. Shin</i>	573
Enhancement of Fermi energy optical emission induced by the band structure in strained-layer InGaAs/InP quantum wells <i>H.A.P. Tudury, F. Iikawa, E. Ribeiro, J.A. Brum, W. Carvalho Jr., A.A. Bernussi, and A.L. Gobbi</i>	575
Band offset determination and excitons in SiGe/Si (001) quantum wells <i>H.H. Cheng, S.T. Yen, and R.J. Nicholas</i>	577
4.5 Nonlinear Optical Studies	
Few-cycle THz spectroscopy of semiconductor quantum structures (invited) <i>K. Unterrainer, R. Bratschitsch, T. Müller, R. Kersting, J.N. Heyman, and G. Strasser</i>	579
Extreme mid-infrared nonlinear optics in semiconductors <i>J. Kono, A.H. Chin, and O.G. Calderon</i>	583
Fine structure of the amplified spontaneous emission of ZnSe laser structures <i>R. Heinecke, U. Neukirch, P. Michler, and J. Gutowski</i>	585
Biexcitonic signatures in femtosecond pulse propagation <i>J. Meinertz, I. Gössling, U. Neukirch, and J. Gutowski</i>	587
Femtosecond response times of large optical nonlinearity in low-temperature grown GaAs/AlAs multiple quantum wells <i>T. Okuno, Y. Masumoto, Y. Sakuma, M. Ito, and H. Okamoto</i>	589
Asymmetric short-period GaAs/AlAs superlattices for observation of low-threshold laser effect <i>V.G. Litovchenko, D.V. Korbutyak, S.G. Krylyuk, A.I. Bercha, H.T. Grahn, and K.H. Ploog</i>	591
Nonlocality induced size-enhancement of excitonic nonlinear response in high quality samples <i>H. Ishihara, K. Cho, K. Akiyama, N. Tomita, Y. Nomura, and T. Isu</i>	593
Recombination mechanism of anti-Stokes photoluminescence in partially ordered GaInP-GaAs heterostructure <i>S.J. Xu, Q. Li, H. Wang, M.H. Xie, S.Y. Tong, and J.R. Dong</i>	595
Spontaneous Raman scattering in GaP-AlGaP heterostructure waveguides <i>T. Saito, K. Suto, M. Kawasaki, T. Kimura, A. Watanabe, and J. Nishizawa</i>	597
Generation mechanisms of coherent phonons in quantum wells revealed <i>K.J. Yee, Y.S. Lim, and D.S. Kim</i>	599

4.6 Carrier Dynamics and Relaxation

Probing and controlling spin-relaxation in GaAs quantum wells (invited) <i>Y. Ohno, R. Terauchi, T. Adachi, F. Matsukura, and H. Ohno</i>	601
Speckle-averaged resonant Rayleigh scattering from quantum well excitons <i>G.R. Hayes, B. Deveaud, V. Savona and S. Haacke</i>	605
Exciton and spin transport by surface acoustic waves in GaAs quantum wells <i>T. Sogawa, P.V. Santos, S.K. Zhang, S. Eshlaghi, A.D. Wieck, and K.H. Ploog</i>	607
Coherent vs. incoherent emission in quantum wells studied by polarisation- and time-resolved spectroscopy <i>G. Aichmayr, L. Viña, S.P. Kennedy, R.T. Phillips, and K. Ploog</i>	609
Ultrafast dynamics of holes in $\text{Si}_{1-x}\text{Ge}_x/\text{Si}$ multiple quantum wells <i>R.A. Kaindl, M. Wurm, K. Reimann, M. Woerner, T. Elsaesser, C. Miesner, K. Brunner, and G. Abstreiter</i>	611
Mechanism of positively charged exciton spin relaxation in CdTe and CdMnTe quantum wells <i>E. Vanelle, P. Kossacki, J. Cibert, T. Amand, P. Renucci, X. Marie, and S. Tatarenko</i>	613
Microscopic simulation of hot-carrier intersubband relaxation in quantum-cascade lasers <i>R.C. Iotti and F. Rossi</i>	615
Reduction of Coulomb scattering in a GaAs/AlGaAs mesoscopic 2DEG disk <i>N. Suzumura, M. Yamaguchi, and N. Sawaki</i>	617
Collective plasma response of interacting electrons localized in disordered GaAs/AlGaAs superlattices <i>Yu.A. Pusep, W. Fortunato, P.P. González-Borrero, A.I. Toropov, and J.C. Galzerani</i>	619
Ultrafast relaxation dynamics of photoexcited carriers in an $\text{In}_{0.53}\text{Ga}_{0.47}\text{As}/\text{InP}$ multiple-quantum well <i>Y. Hamanaka, A. Nakamura, K. Tanase, R. Ohga, Y. Nonogaki, Y. Fujiwara, and Y. Takeda</i>	621
Spin relaxation of negatively charged excitons in CdTe-based quantum wells <i>P. Kossacki, V. Ciulin, M. Kutrowski, J.-D. Ganière, T. Wojtowicz, and B. Deveaud</i>	623
Kinetics of a low density system of indirect excitons in double quantum wells <i>V.I. Yudson</i>	625
Photoexcitation-energy-dependent capture dynamics of excitons in electronically isolated GaAs quantum wells <i>K. Fujiwara, H.T. Grahn, L. Schrottke, and K.H. Ploog</i>	627
Magnetic polaron dynamics at high excitation densities in $\text{Cd}_{1-x}\text{Mn}_x\text{Te}/\text{ZnTe}$ multiple quantum wells <i>R. Pittini, J.X. Shen, M. Takahashi, and Y. Oka</i>	629
Excitonic vs free-carrier spin-relaxation in III-V quantum wells <i>A. Malinowski, P.A. Marsden, R.S. Britton, K. Puech, A.C. Tropper, and R.T. Harley</i>	631
1S-2S-continuum coupling and dephasing in 5 nm GaAs quantum wells <i>A.A. Busch, J.M. Watson, P. Paddon, Z. Wasilewski, and J.F. Young</i>	633
Electric-field-dependent carrier dynamics in an (Al,Ga)As/GaAs double-quantum-well superlattice <i>L. Schrottke, R. Hey, and H.T. Grahn</i>	635
Direct and indirect radiative recombination in strongly excited ZnSe/BeTe superlattices <i>A.A. Maksimov, S.V. Zaitsev, I.I. Tartakovskii, V.D. Kulakovskii, N.A. Gippius, D.R. Yakovlev, W. Ossau, G. Reuscher, A. Waag, and G. Landwehr</i>	637
Radiative recombination and spin relaxation of excitons in $\text{Cd}_{1-x}\text{Mn}_x\text{Te}$ quantum wells <i>M.C. Debnath, J.X. Shen, I. Souma, R. Pittini, and Y. Oka</i>	639
Nonequilibrium electrons in double quantum well structures: Boltzmann equation approach <i>S. Khan-ngern and I.A. Larkin</i>	641
Direct study of spin relaxation processes in 2D ZnCdSe/ZnSe systems through time resolved measurements of polarized exciton emission <i>S. Permogorov, Y. Oka, R. Pittini, J.X. Shen, K. Kayanuma, A. Reznitsky, L. Tenishev, and S. Verbin</i>	643

4.7 Microcavities

Mixed neutral and negatively charged microcavity polaritons (invited) <i>R. Rapaport, E. Cohen, A. Ron, E. Linder, R. de Picciotto, R. Harel, and L.N. Pfeiffer</i>	645
Confined optical modes in photonic molecules and crystals (invited) <i>M. Bayer, A. Forchel, T.L. Reinecke, and P.A. Knipp</i>	649
Non-linear spin polarization dynamics in semiconductor microcavities <i>P. Renucci, X. Marie, T. Amand, M. Paillard, P. Senellart, and J. Bloch</i>	653
Optically pumped quantum dot lasers using high-Q microdisk cavities <i>P. Michler, A. Kiraz, C. Becher, Lidong Zhang, E. Hu, A. Imamoglu, W.V. Schoenfeld, P.M. Petroff</i>	655
Incoherent amplification phenomena in semiconductor microcavities <i>G. Dasbach, T. Baars, M. Bayer, A. Larionov, and A. Forchel</i>	657
Selectively <i>in situ</i> probing of self-assembled InGaAs quantum dots in a planar GaAs microcavity by angle-resolved detection of photoluminescence spectrum <i>J.H. Chen, J.H. Zhao, F.H. Yang, P.H. Tan, J.D. Zhang, Y.P. Zeng, J.Q. You, and H.Z. Zheng</i>	659
Spin quantum beats of exciton-polariton in semiconductor microcavities <i>P. Renucci, X. Marie, T. Amand, M. Paillard, and E. Vanelle</i>	661
Observation of enhanced spontaneous emission coupling factor in blue InGaN microcavities <i>S. Kako, T. Someya, and Y. Arakawa</i>	663
Ballistic transport of exciton-polaritons in a graded quantum microcavity <i>B. Sermage, G. Malpuech, A. Kavokin, and V. Thierry-Mieg</i>	665
Ultrafast polarization switching in a CdTe microcavity <i>M.D. Martín, H. Davies, L. Viña, and R. André</i>	667
Cavity QED of quantum dots embedded in dielectric microspheres <i>H. Wang, X. Fan, and M. Lonergan</i>	669
Polarization of magnetopolaritons in a semiconductor microcavity <i>M.D. Martín, S. Burgas, M. Alonso, L. Viña, F.J. Terán, M. Potemski, and E.E. Mendez</i>	671
Rabi splitting enhancement in planar semiconductor microcavities <i>A. D'Andrea and L. Pilozzi</i>	673
Luminescence properties of CdS quantum dots embedded in monolithic II-VI microcavity <i>T. Tawara, H. Yoshida, H. Kumano, S. Tanaka, and I. Suemune</i>	675
Strongly detuned IV-VI microcavity and microdisk resonances: mode splitting and lasing <i>T. Schwarzl, W. Heiss, G. Springholz, S. Gianordoli, G. Strasser, M. Aigle, and H. Pascher</i>	677
Bottleneck and resonant enhancement in free electron - cavity polaritons scattering in GaAs/AlGaAs microcavities <i>R. Rapaport, A. Qarry, E. Cohen, A. Ron, E. Linder, and L.N. Pfeiffer</i>	679
Biexcitons or bipolaritons in a semiconductor microcavity? <i>P. Borri, W. Langbein, U. Woggon, J.R. Jensen, and J.M. Hvam</i>	681
Theory of microcavity resonant Rayleigh scattering <i>D.M. Whittaker</i>	683
Spatially extended cavity polaritons arising from weakly confined excitons <i>H. Ishihara and J. Kishimoto</i>	685
Stimulation of polariton emission in a homogeneously broadened semiconductor microcavity <i>V. Mizeikis, J. Erland, J.R. Jensen, N.A. Mortensen, and J.M. Hvam</i>	687
Bloch oscillations of light in laterally confined Bragg mirrors and multiple coupled microcavities <i>G. Malpuech, A. Kavokin, A. Di Carlo, P. Lugli, and G. Panzarini</i>	689
Enhanced light transmission through nano-structured surfaces <i>S. Meinlschmidt, R. Windisch, A. Knobloch, P. Kiesel, P. Heremans, and G.H. Döhler</i>	691

Optically pumped lasing at 1.3 μm of GaInNAs-based VCSEL structures <i>M. Hetterich, M.D. Dawson, A.Yu. Egorov, and H. Riechert</i>	693
The interaction between exciton states near the quantum well bandgap and confined photons in GaAs/AlAs microcavities <i>A. Qarry, R. Rapaport, E. Cohen, A. Ron, E. Linder, and L.N. Pfeiffer</i>	695
Resonant Raman scattering in an InAs/GaAs monolayer structure <i>J. Maultzsch, S. Reich, A.R. Goñi, and C. Thomsen</i>	697
Light-exciton coupling in semiconductor microcavities of cylindrical and spherical symmetry <i>R.A. Abram, S. Brand, M.A. Kaliteevski, V.V. Nikolaev, M.V. Maximov, N.N. Ledentsov, C.M. Sotomayor Torres, and A.V. Kavokin</i>	699
4.8 Near-Field Studies	
Near-field spectroscopy of delocalized excitons in single quantum wires <i>F. Intonti, V. Emiliani, Ch. Lienau, T. Elsaesser, R. Nötzel, and K.H. Ploog</i>	701
Scanning near-field optical spectroscopy of buried semiconductor heterostructures <i>M. Hauert, R. Roshan, A.C. Maciel, J. Kim, J.F. Ryan, A. Schwarz, A. Kaluza, Th. Schäpers, and H. Lüth</i>	703
Hints for a Non-thermal distribution of excitons in CdSe/ZnSe quantum islands <i>G. von Freymann, E. Kurtz, C. Klingshörn, Th. Schimmel, and M. Wegener</i>	705
4.9 Intersubband Transitions	
Quantum optics and the observation of electromagnetically induced transparency with QW subbands (invited) <i>C.C. Phillips, E. Paspalakis, G.B. Serapiglia, C. Sirtori, and K.L. Vodopyanov</i>	707
Towards quantum well hot hole lasers <i>P. Kinsler and W.Th. Wenckebach</i>	711
Character of electronic excitations in GaAs-AlGaAs quantum structures <i>E. Ulrichs, C. Steinebach, C. Schüller, Ch. Heyn, W. Hansen, and D. Heitmann</i>	713
Correlation of vertical transport and infrared absorption in GaAs/AlGaAs superlattices <i>M. Helm and G. Strasser</i>	715
Influence of in-plane magnetic field on cyclotron resonance in double-layer two dimensional electron system <i>H. Aikawa, S. Takaoka, A. Kuriyama, K. Oto, K. Murase, S. Shimomura, S. Hiyamizu, T. Jungwirth, and L. Smrčka</i>	717
Circular photogalvanic effect in <i>p</i> -GaAs/AlGaAs MQW <i>S.D. Ganichev, E.L. Ivchenko, H. Ketterl, L.E. Vorobjev, M. Bichler, W. Wegscheider, and W. Prettl</i>	719
Cascading effect in type-II InAs/GaSb/AlSb intersubband light emitter <i>K. Ohtani, H. Sakuma, and H. Ohno</i>	721
Magneto-optical transitions involving a 2DEG confined in Cd(Mn)Te/CdMgTe quantum wells <i>F.J. Teran, M.L. Sadowski, M. Potemski, P. Kossacki, P. Hawrylak, and G. Karczewski</i>	723
Comparison of intersubband relaxation times in GaN/AlGaN and in InGaAs/AlGaAs quantum wells <i>T. Asano, S. Yoshizawa, S. Noda, N. Izuka, K. Kaneko, N. Suzuki, and O. Wada</i>	725
Electronic inelastic light scattering in a periodic δ -doping GaAs multiple quantum well structure <i>C. Kristukat, A.R. Goñi, S. Rutzinger, W. Wegscheider, G. Abstreiter, and C. Thomsen</i>	727
Intersubband electroluminescence using X- Γ carrier injection in a GaAs/AlAs double-quantum-well superlattice <i>C. Domoto, N. Ohtani, K. Kuroyanagi, P.O. Vaccaro, T. Nishimura, H. Takeuchi and M. Nakayama</i>	729
Hot electron optical phenomena in GaAs/AlAs MQW structures in strong lateral electric field <i>L.E. Vorobjev, S.N. Danilov, I.E. Titkov, D.A. Firsov, V.A. Shalygin, A.E. Zhukov, A.R. Koush, V.M. Ustinov, V.Ya. Aleshkin, A.A. Andronov, E.V. Demidov, and Z.F. Krasilnik</i>	731

Novel mid-infrared laser designs based on intraband and interband carrier transitions in quantum wells <i>L.E. Vorobjev, D.A. Firsov, G.G. Zegrya, V.L. Zerova, E. Towe, J.W. Cockburn, and Z.F. Krasil'nik</i>	733
--	-----

5. Heterostructures and Superlattices: Transport

5.1 Transport, Magneto-Transport, and Coherent Effects

Is there a true metallic state in two dimensions? (invited) <i>M.Y. Simmons, A.R. Hamilton, M. Pepper, E.H. Linfield, P.D. Rose, and D.A. Ritchie</i>	735
Self-induced Shapiro effect in semiconductor superlattices <i>F. Löser, M.M. Dignam, Yu.A. Kosevich, K. Köhler, and K. Leo</i>	739
Effects of a parallel magnetic field on the novel metallic behavior in two dimensions <i>K. Eng, X.G. Feng, D. Popović, and S. Washburn</i>	741
Optics with ballistic electrons: Anti-reflection coatings for GaAs-AlGaAs superlattices <i>C. Pacher, G. Strasser, E. Gornik, F. Elsholz, A. Wacker, and E. Schöll</i>	743
Chaotic quantum transport in superlattices <i>T.M. Fromhold, A.A. Krokhin, A.E. Belyaev, C.R. Tench, S. Bujkiewicz, P.B. Wilkinson, F.W. Sheard, L. Eaves, and M. Henini</i>	745
The magnetoresistance of a two-dimensional electron gas in the presence of a spatially random magnetic field <i>A.W. Rushforth, B.L. Gallagher, P.C. Main, C.H. Marrows, B.J. Hickey, E.D. Dahlberg, A.C. Neumann, and M. Henini</i>	747
Giant negative magnetoresistance in two-dimensional antidot arrays: ballistic orbital effect and ballistic weak localization <i>T. Osada, H. Nakamura, and Y. Shiraki</i>	749
Correlation of optical and transport properties of (AlGa)As/GaAs heterostructures <i>L. Gottwaldt, F.J. Ahlers, E.O. Göbel, G. Hein, S. Nau, K. Pierz, and W. Stolz</i>	751
A method of determining potential barrier heights at semiconductor heterointerfaces <i>G.-H. Kim, H.-S. Sim, M.Y. Simmons, D.A. Ritchie, C.-T. Liang, A.C. Churchill, and W.S. Han</i>	753
Conductance quantization in an array of ballistic constrictions <i>S. de Haan, A. Lorke, J.P. Kotthaus, W. Wegscheider, and M. Bichler</i>	755
Transport in asymmetric two-dimensional lateral surface superlattices <i>S. Chowdhury, A.R. Long, J.H. Davies, D.E. Grant, E. Skuras, and C.J. Emeleus</i>	757
Suppression of miniband transport in magnetic field: role of injection process <i>A.A. Krokhin, A.E. Belyaev, T.M. Fromhold, C.R. Tench, H.M. Murphy, S. Bujkiewicz, P.B. Wilkinson, F.W. Sheard, L. Eaves, and M. Henini</i>	759
Quantitative evaluation of electron-electron scattering rate in two-dimensional electron gas by magnetic lateral superlattice <i>M. Kato, A. Endo, M. Sakairi, M. Hara, S. Katsumoto, and Y. Iye</i>	761
Shubnikov-de Haas effect with several filled subbands <i>U. Ekenberg</i>	763
On the origin of beat patterns in the quantum magneto-resistance of gated InAs/GaSb and InAs/AlSb quantum wells <i>A.C.H. Rowe, R.S. Ferguson, and R.A. Stradling</i>	765
Rashba spin splitting in 2D electron and hole systems: Implications for the metal-insulator transition <i>R. Winkler</i>	767
Weak localization in periodically modulated magnetic field <i>Y. Blum, A. Tsukernik, A. Palevski, T.A. Shutenko, B.L. Altshuler, I.L. Aleiner, A. Rudra, and E. Kapon</i>	769

Intrinsic mobility limits in polarization induced two-dimensional electron gases <i>D. Jena, Y.P. Smorchkova, C.R. Elsass, A.C. Gossard, and U.K. Mishra</i>	771
Combined <i>S</i> - and <i>Z</i> -shaped current bistability induced by charging of quantum dots <i>A.E. Belyaev, L. Eaves, S.A. Vitusevich, P.C. Main, M. Henini, A. Förster, N. Klein, and S.V. Danylyuk</i>	778
Interaction of μ with spin current in GaAs/GaAsP/Si <i>E. Torikai, Y. Ikedo, A. Ihori, K. Shimomura, K. Nagamine, T. Saka, and T. Kato</i>	775
Metallic behaviour and temperature dependent screening in p-SiGe <i>V. Senz, T. Ihn, T. Heinzel, K. Ensslin, G. Dehlinger, U. Gennser, and D. Gruetzmacher</i>	777
Anisotropic magneto-transport properties of 70 nm-period lateral surface superlattices in high magnetic fields <i>M. Akabori, J. Motohisa, and T. Fukui</i>	779
Thermopower of a two-dimensional antidot lattice <i>A.G. Pogosov, M.V. Budantsev, A.E. Plotnikov, A.K. Bakarov, and A.I. Toropov</i>	781
Magnetotransport in two-dimensional lateral superlattices in strongly coupled electron-hole gases <i>B. Kardynal, R.J. Nicholas, J. Rehman, K. Takashina, and N.J. Mason</i>	783
Semiclassical origin of the 2D metallic state in high mobility Si-MOS and Si/SiGe structures <i>G. Brunthaler, A. Prinz, G. Pillwein, G. Bauer, K. Brunner, G. Abstreiter, T. Dietl, and V.M. Pudalov</i>	785
Zero-bias conductance anomaly in GaAs/AlGaAs modulation doped field-effect transistors <i>S. Skaberna, U. Kunze, D. Reuter, and A.D. Wieck</i>	787
Magnetotransport and capacitance investigations of strongly coupled but spatially separated 2D electron and hole systems <i>M. Pohl, M. Lynass, W. Dietsche, K. von Klitzing, K. Eberl, and R. Mühle</i>	789
First principles study of spin-electronics: Zero-field spin-splitting in superlattices <i>J.A. Majewski, P. Vogl, and P. Lugli</i>	791
A comparative study of 'metallic' and 'insulating' behaviour of the two-dimensional electron gas on (100) and vicinal surfaces of Si MOSFET <i>S.H. Roshko, S.S. Safonov, A.K. Savchenko, A.G. Pogosov, and Z.D. Kvon</i>	793
The role of surface-localized states in the in-plane transport properties of superlattices <i>A.B. Henriques</i>	795
High-mobility heterostructure as a new kind of chaotic billiards <i>L.D. Shvartsman</i>	797
Gate-controlled very large spontaneous spin-splittings in normal $\text{In}_{0.75}\text{Ga}_{0.25}\text{As}/\text{In}_{0.75}\text{Al}_{0.25}\text{As}$ heterojunctions grown on GaAs substrates <i>S. Yamada, Y. Sato, S. Gozu, and T. Kikutani</i>	799
Field domains in semiconductor superlattices: Dynamic scenarios of multistable switching <i>A. Amann, A. Wacker, L.L. Bonilla, and E. Schöll</i>	801
Angular dependent magnetoresistance oscillations in semiconductor superlattice with incoherent interlayer coupling <i>M. Kuraguchi, E. Ohmichi, T. Osada, and Y. Shiraki</i>	803
Optical and transport properties of modulation doped InAs/GaAs superlattices <i>V.A. Kulbachinskii, R.A. Lunin, V.G. Kytin, A.V. Golikov, V.A. Rogozin, V.G. Mokerov, Yu.V. Fedorov, and A.V. Hook</i>	805
Weak localization effects in a wide parabolic quantum well <i>N.M. Sotomayor, G.M. Gusev, J.R. Leite, N.T. Moshegov, and A.I. Toropov</i>	807
Magnetotransport properties of multisubband semiconductor structures <i>N.S. Averkiev, L.E. Golub, S.A. Tarasenko, and M. Willander</i>	809

Effect of buffer layer thickness on improvement of modulation doped CdTe/CdMgTe heterostructures grown on GaAs substrate <i>D. Wasik, M. Baj, L. Dmowski, J. Siwiec-Matuszyk, J. Przybytek, E. Janik, T. Wojtowicz, and G. Karczewski</i>	811
Domain formation in a one-dimensional superlattice with phonon scattering <i>L.G. Mouroukh, A.Yu. Smirnov, N.J.M. Horing, and V.I. Gavrilenko</i>	813
Asymmetric carrier diffusion and phonon-wind-driven transport in an InGaAs-InP quantum well <i>A.F.G. Monte, S.W. da Silva, P.C. Morais, J.M.R. Cruz, and A.S. Chaves</i>	815
5.2 Magnetic Superlattices	
Nonzero Hall resistance in a spatially fluctuating magnetic field with zero mean <i>A.A. Bykov, G.M. Gusev, J.R. Leite, A.K. Bakarov, N.T. Moshegov, D.K. Maude, M. Casse, and J.C. Portal</i>	817
5.3 Tunneling and Resonant Tunneling	
Optical detection of ballistically injected electrons in III/V heterostructures <i>M. Kemmerink, K. Sauthoff, P.M. Koenraad, J.W. Gerritsen, H. van Kempen, and J.H. Wolter</i>	819
Electron tunneling time between double quantum dot <i>A. Tackeuchi, Y. Nakata, T. Kuroda, K. Mase, and N. Yokoyama</i>	821
Direct measurement of the AlAs X-band Fermi surface <i>H.-S. Im, L.E. Bremme, P.C. Klipstein, A.V. Kornilov, R. Grey, and G. Hill</i>	823
Miniband transport and Stark-cyclotron-resonance in InAs/GaSb superlattices <i>V.J. Hales, R.J. Nicholas, and N.J. Mason</i>	825
Effective mass anisotropy of Γ -electrons in GaAs/AlGaAs quantum wells due to interface band mixing <i>T. Reker, H. Im, H. Choi, L.E. Bremme, Y.C. Chung, R. Grey, G. Hill, and P.C. Klipstein</i>	827
Vertical transport and interband luminescence in type II InAs/GaSb/InAs heterostructures <i>M. Roberts, N.J. Mason, S.G. Lyapin, Y.C. Chung, and P.C. Klipstein</i>	829
Resonant tunneling of holes under in-plane uniaxial stress <i>Y.C. Chung, T. Reker, L.E. Bremme, R. Grey, and P.C. Klipstein</i>	831
Resonant tunneling through zero-dimensional impurity states: Effects of a finite temperature <i>P. König, U. Zeitler, J. Königmann, T. Schmidt, and R.J. Haug</i>	833
Temperature dependence of resonant tunneling characteristics in a p-type GaAs/AlAs double-barrier structure <i>M. Ono, N. Nishioka, M. Morifuji, and C. Hamaguchi</i>	835
Negative differential resistance and current self-oscillation in doped GaAs/AlAs superlattices <i>J.N. Wang, C.Y. Li, X.R. Wang, B.Q. Sun, Y.Q. Wang, W.K. Ge, D.S. Jiang, and Y.P. Zeng</i>	837
Photocurrent self-oscillations in weakly coupled, type-II GaAs/AlAs superlattices embedded in p-i-n and n-i-n diodes <i>N. Ohtani, M. Rogozia, C. Domoto, T. Nishimura, and H.T. Grahn</i>	839
Zener-phonon resonances in the quantum transport of multiband semiconductor superlattices <i>P. Kleinert</i>	841
Correlation between a remote electron and a two-dimensional electron gas in resonant tunneling devices <i>H. Kato and F.M. Peeters</i>	843
Spin effects in InAs quantum dots: Tunneling experiments in tilted magnetic fields <i>J.M. Meyer, I. Hapke-Wurst, U. Zeitler, R.J. Haug, H. Frahm, A.G.M. Jansen, and K. Pierz</i>	845
Carrier transport affected by hole-subband resonances in a strained GaAs/InAlAs superlattice <i>M. Hosoda, K. Kuroyanagi, N. Ohtani, and T. Aida</i>	847
Disorder-enhanced tunneling transport through doping barriers <i>R. Elpelt, O. Wolst, H. Willenberg, S. Malzer, and G.H. Döhler</i>	849

Response time of the double-barrier heterostructures with resonant tunneling <i>M.N. Feiginov</i>	851
Current self-oscillations with discrete frequencies in weakly coupled semiconductor superlattices <i>M. Rogozia, H.T. Grahn, and R. Hey</i>	853
Observation of the scattered electrons in the resonant tunneling regime using a three-terminal quantum-well heterostructure <i>G.-G. Kim, D.-W. Roh, S.-W. Paek, K.-M. Koh, K.E. Pyun, and C.-H. Kim</i>	855
5.4 Hopping, Localization	
Nature of the localized phase in a two-dimensional electron system <i>I.E. Itskevich, R.J.A. Hill, S.T. Stoddart, H.M. Murphy, A.S.G. Thornton, P.C. Main, L. Eaves, M. Henini, D.K. Maude, and J.C. Portal</i>	857
Evidence for screening breakdown near the metal-to-insulator transition in two dimensions <i>W. Jantsch, Z. Wilamowski, N. Sandersfeld, and F. Schäffer</i>	859
Coulomb interaction and density of states in amorphous $\text{Si}_x\text{Ge}_{1-x}$ films <i>K. Nakada, K. Nara, N. Aoki, and Y. Ochiai</i>	861
Studies of localization in an interacting two-electron system <i>J. Talamantes, M. Pollak, and I. Varga</i>	863
5.5 Phonons, Plasmons	
Bridging the gap with cleaved edge overgrowth superlattices : on minigaps, magnetic breakdown and quantum interference in artificial bandstructures (invited) <i>R.A. Deutschmann, W. Wegscheider, C. Albrecht, J.H. Smet, M. Rother, M. Bichler, and G. Abstreiter</i>	865
Electron wires driven by a surface acoustic wave and nonlinear acoustoelectric interactions in quantum wells <i>A.O. Govorov, A.V. Kalameitsev, V.M. Kovalev, H.-J. Kutschera, M. Streibl, M. Rotter, and A. Wixforth</i>	869
Generation and detection of picosecond acoustic phonon pulses in a double quantum well structure <i>I. Ishii, O. Matsuda, T. Fukui, J.J. Baumberg, and O.B. Wright</i>	871
Amplification and generation of high-frequency coherent acoustic phonons under the drift of 2D-electrons <i>M.A. Stroscio, S.M. Komirenko, K.W. Kim, A.A. Demidenko, and V.A. Kochelap</i>	873
Observation of a two-dimensional plasmon in a metallic monolayer on silicon surface <i>T. Nagao, T. Hildebrandt, M. Henzler, and S. Hasegawa</i>	875
First- and second-order Raman spectroscopy of $^{70}\text{Ge}_n/^{76}\text{Ge}_n$ isotope superlattices <i>K. Morita, K.M. Itoh, M. Nakajima, H. Harima, K. Mizoguchi, Y. Shiraki, and E.E. Haller</i>	877
Raman scattering in Ge quantum dot superlattices <i>A. Milekhin, N. Stepina, A. Yakimov, A. Nikiforov, S. Schulze, T. Kampen, and D.R.T. Zahn</i>	879
Plasmon-phonon coupled mode excitation in a hot 2DEG observed by a phonon pulse technique <i>J.K. Wigmore, H.A. Al Jawhari, A.G. Kozorezov, and M. Sahraoui-Tahar</i>	881
Peculiarities of phonons in strained short-period GaN/AlN superlattices: A <i>first-principles</i> study <i>J.-M. Wagner and F. Bechstedt</i>	883
5.6 High-Frequency Transport, Microwaves, Noise, Short Pulses	
Study of ambipolar diffusion and drift of spatially separated charge carriers <i>M. Beck, M. Vitzethum, D. Streb, P. Kiesel, C. Metzner, S. Malzer, and G.H. Döhler</i>	885
Localization of carriers and Bloch oscillations in quantum dot superlattices in dc electric field <i>R.A. Suris and I.A. Dmitriev</i>	887
A novel quantum transport mechanism in biased quantum-box superlattices under terahertz irradiation <i>P. Kleinert and V.V. Bryksin</i>	889
On the spectra of field and current oscillations due to laser irradiation in superlattices <i>Yu.A. Romanov and Ju. Yu. Romanova</i>	891